

An Ectopic Intrathyroidal Parathyroid Adenoma in a Minimally Symptomatic Hypercalcemic Patient: A Rare Case Presentation

Hinal Rathi^{1*}, Patrick A. Dolan² and John Webb²

¹Medical Student, Arkansas College of Osteopathic Medicine, Fort Smith, Arkansas, USA

²Department of General Surgery, CHI St. Vincent Hospital, Hot Springs, Arkansas, USA

*Corresponding author: Hinal Rathi, Medical Student, Arkansas College of Osteopathic Medicine, Fort Smith, Arkansas, USA.

E-mail: hrathi@arcomedu.org

Received: May 25, 2022; Accepted: June 06, 2022; Published: June 17, 2022

Abstract

A common etiology of hypercalcemia is Primary Hyperparathyroidism. Eighty percent of the time a single adenoma is present. Labs usually show an elevated Parathyroid hormone. Head and neck US, Sestamibi and CT scanning are important for localization. In our case, diagnostic results suggested a right lower pole location for a parathyroid adenoma. Referral to General Surgery was made for focused neck dissection. Bilateral neck exploration was required as only three glands were identified. After exploration of possible ectopic inferior gland locations, a right thyroid lobectomy was performed. This ectopic location was confirmed via intraoperative serial PTH measurements. This case illustrates a rare ectopic location for a parathyroid adenoma.

Keywords: Hypercalcemia; Primary hyperparathyroidism; Ectopic parathyroid adenomas

Introduction

Primary Hyperparathyroidism (PHP) is a disease of abnormal calcium and phosphate homeostasis. The etiology being an elevated Parathyroid hormone (PTH) level with subsequent hypercalcemia and low phosphate levels [4]. The increase in serum calcium levels are related to increased bone resorption and increased renal and gut uptake. It is the 3rd most common endocrine disorder.

When hypercalcemia is present, a PHP etiology is usually considered with a twice normal or greater intact PTH level. Risk factors for PHP include hereditary causes such as multiple endocrine neoplasias, exposure to ionizing radiation, prolonged lithium therapy, reduced physical activity, and low calcium intake [4,6]. At times, patients are mildly symptomatic with nonspecific symptoms such as fatigue, mild depression, or cognitive impairment [5].

However, symptomatic patients with PHP may present with multi-organ disorders such as nephrolithiasis, renal insufficiency, peptic ulcer disease, pancreatitis, bone pain, secondary fractures, osteoporosis, osteitis fibrosa cystica, lethargy, psychosis or even coma with severe hypercalcemia [5]. Also, with severe hypercalcemia with a parathyroid etiology, carcinoma should be considered.

Case Description

This case study describes a 46-year-old, mildly symptomatic, African American male patient who was found to have, through routine lab testing through his primary care physician, a calcium of 10.8 mg/dL (normal 8.6 - 10.3 mg/dL). Subsequent labs revealed a PTH level of 191.5 pg/mL (normal 14-65 pg/mL), and a vitamin D level of 29 ng/mL (normal 20 ng/mL - 50 ng/mL). Calcium that is bound to vitamin D is an inactive form.

The patient's past medical history was significant for gout. However, the patient denied a history of head or neck radiation, kidney stones, peptic ulcer disease, or a previous malignancy. Review of systems was positive for fatigue, but negative for myalgias and malaise. Family history was negative for endocrine neoplasias.

The Endocrinologist rechecked lab work multiple times which showed a PTH level in the range of 400- 500 pg/mL and a Ca level in the range of 10.8-13.5 mg/dL. Ultrasound of the neck showed a lower right inferior mixed solid/cystic appearing dominant thyroid nodule with small punctate calcifications. FNA showed benign fibroadipose tissue containing heterotopic thymic tissue with cystic change admixed with two microscopic foci of benign thyroid tissue. Lymph node tissue fragments showed no evidence of metastatic disease. (Figure 1). It should be noted that Pathology reexamined the FNA slides post-surgery and a single slide suggested parathyroid tissue. Finally, a Sestamibi scan showed evidence of a parathyroid adenoma at the level of the inferior right thyroid gland (Figure 2).




Figure 1: Head and Neck Ultrasound Findings: The right lobe is enlarged and mostly homogenous and measures 3.22 x 2.37 x >6.34 cm in transverse x anterior-posterior x longitudinal dimensions. Within the right lobe, there is a superficial anechoic cystic-appearing nodule measuring 0.96 x 0.69 x 1.09 cm and an inferior large mixed solid/cystic appearing anechoic/isoechoic heterogeneous mass measuring 2.06 x 3.61 x 3.28 cm.




Figure 2: Sestamibi Scan Findings: There is diffuse uptake of tracer noted in the salivary glands and thyroid. Mild increased uptake is noted in the inferior right thyroid. On the three-hour delayed imaging, focal radiotracer persists at the level of the inferior right thyroid, concerning for a parathyroid adenoma. The activity washes out of the remainder of the thyroid gland. Findings concerning for a parathyroid adenoma at the level of the inferior right thyroid gland.

The patient was referred to General Surgery for a focused right parathyroidectomy. Intraoperatively, bilateral neck exploration was required as only three of the four parathyroid glands were initially identified. After an extensive exploration of possible ectopic locations for an inferior gland, a right thyroid lobectomy was performed. The presence of an intrathyroid adenoma was confirmed through intraoperative serial PTH measurements. Pre-operative PTH level was 354 pg/mL; 10-min post gland removal PTH level was 10 pg/mL and 15-min post gland removal PTH level was 12 pg/mL. Post-op serial sections of the tissue specimen showed a thyroid lobe containing an intrathyroidal parathyroid adenoma (4 cm), along with an incidental benign colloid nodule (0.6 cm).

Discussion

For cases of hypercalcemia due to primary hyperparathyroidism, a single adenoma is present 80% to 85% of the time [5]. Neck ultrasound and Sestamibi scanning are used to visualize the location of the parathyroid adenoma [4,5]. If ultrasound results are unclear as to location, CT scanning is often used. Parathyroid adenomas are more common in those greater than 50 years of age and women more than men [7]. Parathyroidectomy is usually recommended in cases of less than 50 years of age, history of kidney stones, serum calcium level greater than 1mg/dL above the upper limit of normal, and reduced bone density [2]. However, surgery can also be offered to asymptomatic and minimally symptomatic elderly patients under certain circumstances [6].

Normal parathyroid gland anatomy consists of four glands, two superior and two inferior [2,5]. During embryogenesis the fourth branchial pouch gives rise to the superior glands and the third branchial pouch gives rise to the inferior glands and the thymus [2,5]. Parathyroid adenomas can be found in several ectopic locations due to the defects caused during the embryological descent of the parathyroid glands [2].

Ectopic superior parathyroid glands are commonly found in the tracheoesophageal groove and the retroesophageal region. Ectopic inferior parathyroid glands can be found in the anterior mediastinum, usually associated with the thymus, within the thyroid gland, and within the carotid sheath. [5]. The location of the inferior parathyroid gland is more variable as it takes the longer migratory route during embryogenesis [5]. In one of the clinical studies, 202 patients were analyzed for ectopic locations of parathyroid glands: 38% percent were found to be in the thymus, 31% in the retroesophageal region and 18% intrathyroidal [1]. The intrathyroidal location being the most rare.

Preoperative localization of the adenoma is very important. Imaging modalities such as dual phase ^{99m}Tc- Sestamibi scintigraphy is currently the gold standard in diagnosing and localizing parathyroid adenomas. FNA can not only assist in localizing ectopic adenomas, but also suggest the presence of a concomitant thyroid carcinoma in a prominent thyroid nodule. In this case, Ultrasound imaging and Sestamibi scanning suggested a right lower lobe location. FNA was used to possibly obtain a tissue diagnosis preoperatively. Initially however, an intrathyroid ectopic location was not confirmed by FNA. Serial PTH measurements were thereafter utilized perioperatively which assisted in confirming that the adenoma had been excised.

Conclusion

Parathyroid adenomas are primarily treated through surgical resection with a curative rate in 95- 97% of cases [1,3]. Generally, 6-16% of parathyroid adenomas occur in ectopic locations. Where the inferior glands are concerned however, the mediastinum and intrathyroid locations are most likely. This is due to abnormal migration during embryogenesis [2]. Occasionally, a fourth, usually ectopic gland cannot be found resulting in continued refractory hypercalcemia postoperatively. Therefore, thorough perioperative evaluation is necessary to avoid post-surgical failure. Finally, to avoid leaving an ectopic gland, it is important to have an experienced surgeon conduct a methodical intraoperative search. When only three glands are found and the missing gland is in the inferior position, the final step is a thyroid lobectomy on the side of the missing gland.

REFERENCES

1. Roy M, Mazeh H, Chen H, et al. Incidence and localization of ectopic parathyroid adenomas in previously unexplored patients. *World J Surg.* 2013; 37: 102-106.
2. Noussios G, Anagnostis P, Natsis K. Ectopic parathyroid glands and their anatomical, clinical and surgical implications. *Exp Clin Endocrinol Diabetes.* 2012; 120: 604-610.
3. Eslamy HK, Ziessman HA. Parathyroid scintigraphy in patients with primary hyperparathyroidism: 99mTc sestamibi SPECT and SPECT/CT. *Radiographics.* 2008; 28: 1461-1476.
4. Wilhelm SM, Wang TS, Ruan DT, et al. The American Association of Endocrine Surgeons Guidelines for Definitive Management of Primary Hyperparathyroidism. *JAMA Surg.* 2016; 151: 959-968.
5. Guilmette J, Sadow PM. Parathyroid Pathology. *Surg Pathol Clin.* 2019; 12: 1007-1019.
6. Khan A, Hanley D, Rizzoli R, et al. Primary hyperparathyroidism: review and recommendations on evaluation, diagnosis, and management. A Canadian and international consensus. *Osteoporosis International.* 2016; 28: 01-19.
7. Edafe O, Collins EE, Ubhi CS, et al. Current predictive models do not accurately differentiate between single and multi-gland disease in primary hyperparathyroidism: a retrospective cohort study of two endocrine surgery units. *Annals of the Royal College of Surgeons of England.* 2018; 100: 140-145.